

OFFICE MEMORANDUM

Subject: Inviting Comments/Suggestions on the Draft “Guidelines/SOP for Blood Transfusion & Blood Bank for Animals in India” reg.

The Department of Animal Husbandry and Dairying (DAHD), Ministry of Fisheries, Animal Husbandry and Dairying, Government of India, has developed a draft document titled **“Guidelines/Standard Operating Procedure (SOP) for Blood Transfusion & Blood Bank for Animals in India”** aimed at standardizing and facilitating the safe use of blood transfusion practices in veterinary care.


2. All concerned institutions, experts, and members of the public are hereby requested to examine the attached draft documents and submit their valuable comments, suggestions, and feedback to the Department for further refinement and finalization of the guidelines.

3. The comments/suggestions may be sent via email to the following IDs:

- sharma.aruna@gov.in
- adhiraj.lolh@gov.in

4. The inputs/feedback may kindly be submitted within **10 (ten) days** from the date of issuance of this Office Memorandum.

Encl.: Draft “Guidelines/SOP for Blood Transfusion & Blood Bank for Animals in India and also hosted in the DAHD website i.e . dahd.gov.in.


(Dr. Adhiraj Mishra)


Assistant Commissioner (AH)

Distribution

1. Director General Health Services, Govt. of India, New Delhi
2. Deputy Director General (Animal Sciences), ICAR, New Delhi
3. Director, Animal Husbandry all States/UTs
4. Vice Chancellors of all Veterinary Universities
5. Dean of all Veterinary Colleges of India
6. Secretary, Veterinary Council of India
7. Secretary, Animal Welfare Board of India
8. HoD, NIC, Krishi Bhawan for uploading in the Department website.

Copy to

1. Sr PPS to Secretary, AHD
2. PPS to AHC
3. PPS to AS(LH)
4. PPS to JS(NLM)



**Department of Animal Husbandry & Dairying
Ministry of Fisheries Animal Husbandry & Dairying
Government of India
July 2025**

Executive Summary

India's livestock and companion animal sectors play a pivotal role in the country's agricultural economy and rural livelihoods. However, emergency and critical care veterinary services, particularly blood transfusion and blood banking, remain largely unstructured and presently lacks comprehensive national protocols governing veterinary blood transfusion.

This document outlines the Standard Operating Procedure (SOP) for blood transfusion and blood banking services for companion animals and livestock in India. It establishes national guidelines for donor selection, blood collection, processing, storage, transfusion protocols, donor welfare, and legal and ethical standards. The SOP aims to standardize transfusion practices, ensure donor animal welfare, and integrate One Health principles to minimize zoonotic risks.

Key highlights include recommendations for the establishment of state-regulated blood banks, mandatory donor health screenings, standard storage conditions for blood components, clear transfusion monitoring protocols, and a robust legal framework to support nationwide implementation. Emphasizing ethical considerations, voluntary non-remunerated donations, and public awareness initiatives, this document also advocates for capacity building through veterinary training programs and future research in advanced transfusion technologies.

This SOP provides a policy and procedural foundation to establish a structured and ethical animal blood banking network across India, contributing significantly to veterinary healthcare and animal welfare.

Table of Contents

1. Introduction
2. Objectives
3. Scope and Definitions
4. Indications for Blood Transfusion
5. Donor Selection and Welfare
6. Blood Collection and Processing
7. Component Separation and Storage
8. Transfusion Protocols
9. Monitoring and Reaction Management
10. Establishment of Blood Banks
11. Legal and Regulatory Framework
12. Ethical Considerations and Owner Education
13. Training and Curriculum Development
14. Future Directions
15. Reference

1. Introduction

India's livestock and companion animal sector is among the most diverse and economically significant in the world. With over **537 million livestock** and more than **125 million companion animals**, the country hosts the **world's largest population of cattle and buffalo**, the **second-largest goat population**, and a rapidly growing number of dogs and cats. The animal husbandry sector contributes approximately **5.5% to the national GDP** and **30% to the agricultural GDP**, serving as a backbone for food security, rural livelihoods, and public health.

With the evolution of veterinary practice in India, including advanced diagnostics and therapeutic capabilities, there is an urgent need to formalize **critical and emergency veterinary care services**, including **blood transfusion support across species**. Blood transfusion is increasingly recognized globally as a life-saving intervention, essential for managing trauma, severe anemia, surgical blood loss, infectious diseases, and coagulation disorders.

At present, India lacks a **national regulatory framework** and **standardized protocols** guiding veterinary transfusion medicine with structured blood banking and transfusion practices. In India, most animal blood transfusions are still performed in emergency settings, relying on hospital-available or client-owned donors, without consistent screening, blood typing, or standard operating protocols.

This SoP aims to address these gaps by providing structured, ethical, and scientifically sound guidance on donor selection, blood collection, processing, storage, transfusion procedures, and safety monitoring. It is developed to support veterinary practitioners, institutions, and policymakers in establishing and managing blood banks for both **companion animals** and **livestock species**, while ensuring animal welfare and public confidence.

2. Objectives

- Establish national guidelines for donor selection, blood collection, processing, storage, and transfusion.
- Ensure donor animal welfare and adherence to ethical standards.
- Standardize transfusion protocols to optimize clinical outcomes.
- Ensure traceability and safety across the transfusion chain.
- Integrate One Health principles to manage zoonotic disease risks.
- Lay the foundation for a national blood bank network for animals
- Promote awareness and impart training among veterinary professionals

3. Scope and Definitions

This document applies to veterinary blood transfusion of companion animals (dogs, cats) and farm animals (cattle, buffalo, sheep, goats) across India in case of emergency, critical and planned care.

Key Definitions:(more points to be added)

- Animal: Domesticated mammals under veterinary care
- Community Sourced Donor: Animal voluntarily donating blood with owner consent
- Donor Welfare: Protection of donor animals from harm, distress, or exploitation
- Blood Components: Whole blood, RBC, plasma, platelets, cryoprecipitate
- Adverse Event: Any unexpected reaction during or after donation or transfusion

4. Indications for Blood Transfusion

The various components of blood can be used in different conditions as indicated below:

4.1 Fresh Whole Blood

Fresh Whole Blood that is less than 8 hours old and has not been refrigerated and has all cellular and fluid components of blood. Stored Whole Blood is blood that is more than 8 hrs old. It can be stored at 33.8-42.8 °F (1 - 6 °C) for a maximum of 28-35 days.

- a. Treatment for anemia with thrombopathy or severe thrombocytopenia and disseminated intravascular coagulation
- b. Animals with severe trauma requiring massive transfusion
- c. Animals suffering from blood-related protozoal diseases like Anaplasmosis, Babesiosis
- d. Animals suffering from anaemia with hypoproteinemia
- e. For intravascular volume expansion and oxygen support

4.2 Packed RBCs

Contains concentrated RBCs with a PCV of 80 percent. Depending on the anticoagulant used, it can be stored for 28 – 35 days.

- a. Animals with symptomatic anemia including anemia due to hemorrhage, hemolysis, renal disease and decreased production of RBCs (bone marrow suppression)
- b. Helps to increase RBC concentration in animals that require oxygen support

4.3 Platelet Rich Plasma (Platelet concentrate)

Obtained by centrifuging fresh blood at a rate slower than that used for production of packed cell volume and plasma. Difficult to store. Can be used within 48 hrs if kept away from light at room temperature. Should not be refrigerated.

- a. Consistently administered to animals with life threatening hemorrhage due to marked thrombocytopenia

b. To animals with hemorrhage secondary to congenital or acquired thrombopathy

5. Blood and Blood Volumes of Animals

Blood is the vascular connective tissue composed of fluid parts termed plasma and of corpuscles viz - erythrocytes or red blood cells, leucocytes or white blood cells and platelets.

The shape and size of red blood cells vary in different animals eg. Dog's RBC is a typical biconcave disk, whereas the Goat's RBC is more spherical. RBC shape in Camel is elliptical, in Deer it is sickle shaped. Humans and animals have blood volumes of 7% to 9% of their body weight. When there is a reduction in the blood volume, transfusion is generally required.

Animals	Blood volume (ml/kg)	Average body wt (kg)	Total Blood Volume (ml)
Cattle	55	300	16500
Goat	66	20	1320
Sheep	66	35	2310
Dog	86	25	2150
Cat	55	5	275
Pig	65	60	3900
Horse	76	450	34200
Monkey (Rhesus)	54	8	432
Rabbit	56	4	224
Guinea pig	75	1.2	90
Ferret	75	2	150

5.1 Blood Types

Every species has specific cell surface antigens on its RBC which differ leading to the development of the blood groups in animals. In animals, there is a combination of blood factors leading to the terminology of Blood grouping system rather than blood groups.

Blood groups are produced by the presence of species-specific antigens on the cell membrane of red blood cells. These factors cause a devastating role in causing adverse-systemic reactions following transfusion. Antigens can also get adsorbed onto the leucocytes, platelets, and plasma protein failing transfusion. When these antigens encounter other organisms either by transfusion or through transplacental exposure, it induces the production of antibodies in the young ones and even causes neonatal isoerythrolysis in certain species like dog.

Species	No. of blood group	Description
Cattle	11	A, B, C, F, J, L, M, R, S, T, Z
Sheep	7	A, B, C, D, M, R, X
Goat	5	A, B, C, M, J
Horse	8	A, C, D, K, P, Q, U, T
Pig	11	A-O, C, F, G, H, I, J, K, L, N, O
	15	
Dog	9	DEA1.1,1.2,3,4,5,6,7,8, Dal antigen
Cat	4	A, B, AB, Mik antigen
Chicken	13	A, B, C, D, E, H, I, J, K, L, N, P, R

6. Mandatory Blood Typing and Cross-Matching:

6.1 Blood Typing: It is a technique by which the surface antigen of the erythrocytes can be identified. It works well to screen the blood donors of the animals before crossmatching and transfusion. They are determined in the laboratory and even blood-typing kits like card-based agglutination assay, immunochromatographic cartridge and gel column diffuse assay are available in the market. These cards are available for both felines and canine's blood-typing and can detect DEA 1.1 and felines type A, B, and AB. In samples of canine blood that is potentially DEA 1.1 – negative, the laboratory can also type the blood or DEA 1.2 to confirm that RBCs are negative for both DEA 1 alleles which could mean that dog is a Universal donor

6.2 Cross-Matching: Ensures compatibility between donor and recipient blood to prevent transfusion reactions. A major, minor, and auto control crossmatch should be performed although the minor crossmatch is rarely used in dogs. The major crossmatch should always be compatible at room temperature and at 37°C. The end reaction to being noticed is hemolysis and agglutination

Major Cross-Match: Tests donor red blood cells against recipient plasma to detect antibodies against donor antigens. To perform the test, mix 2 drops of recipient plasma with 1 drop of donor RBC suspension. Incubate the mixture for 30 minutes at 37°C, then centrifuge at 1500 rpm for 2 minutes. After centrifugation, examine the sample both microscopically for agglutination and macroscopically for hemolysis. Visible hemagglutination indicates incompatibility.

Major cross matching = Recipient's Plasma + Donor's RBC

Minor Cross-Match: Tests donor plasma against recipient red blood cells, less common as most transfusions use packed red cells. To carry out the test, mix 2 drops of donor plasma

with 1 drop of recipient RBC suspension. Incubate the mixture for 30 minutes at 37°C, then centrifuge at 1500 rpm for 2 minutes. Following centrifugation, examine the sample microscopically for agglutination and macroscopically for hemolysis.

Minor cross matching = Recipient's RBC + Donor's Plasma

Recipient's control = Recipient's Plasma + Recipient's RBC

There are several in-house cross-matching tests like Alvedia, Rapid Vet H Major for both feline and canine species. Cross matching is not required before a first transfusion in a dog, due to the lack of naturally occurring alloantibodies (NOAbs), but is strongly recommended > 4 days after a subsequent transfusion. Due to the presence of highly immunogenic NOAbs in cats, major cross matching is strongly recommended both before a first transfusion in cats and then > 2 days after any subsequent transfusion.

6.3 Veterinary Blood Bags Availability

Human Blood Bags Availability

100 ml CPDA / 350 ml CPDA / 300 ml transfer bags

6.4 Blood Collection Equipment

- Blood collection bags (with CPDA-1 or CPD anticoagulant) – single, double, triple or quadruple
- Needles – 16G or 18G sterile blood collection needles
- Blood bag weighing scale/mixer – to prevent clotting and ensure volume accuracy
- Sphygmomanometer cuff / Pressure bag – helps collect blood from jugular vein by mild pressure if needed
- Labeling tools – permanent markers, barcodes, stickers for unit ID and data
- Sterile gloves, antiseptic swabs, surgical clippers

6.5 Donor Selection and Screening

To ensure safe and effective veterinary blood transfusion, donor animals—whether canine, feline, or livestock—must meet the following criteria:

A. General Health and Suitability

- Must be **clinically healthy** with no signs of systemic illness.
- Free from **tick-borne and vector-borne diseases**.
- Should be **docile and temperamentally suitable** for handling during donation.
- Not under **any ongoing medication or treatment**.

B. Age and Weight Requirements

- **Dogs:** Age **1–8 years**, minimum body weight **25 kg**.
- **Cats:** Age ideally **1–5 years**, minimum body weight **4 kg**, not obese.
- **Livestock:** Healthy adults selected based on species-specific clinical norms.

C. Vaccination and Reproductive Health

- Donors must be **fully vaccinated**, especially against **rabies**.
- Should be **regularly dewormed**.
- **Female animals** must not be **pregnant** or **recently lactating**.
- A **minimum 30-day interval** between successive donations is mandatory.

D. Donation Frequency

- **Dogs:** Eligible for donation every **4–6 weeks**.
- **Cats:** Eligible every **8–12 weeks**.

E. Mandatory Pre-Donation Screening

- A **clinical examination** and **medical history review** must be conducted before every donation.
- **Laboratory parameters** to be assessed:
 - **Packed Cell Volume (PCV):** $\geq 35\%$
 - **Hemoglobin (Hb):** ≥ 10 g/dL
 - **Platelet count:** $> 200,000/\mu\text{L}$
 - **Total protein:** Within species-specific normal limits

F. Infectious Disease Screening

Donors must test **negative for transfusion-transmissible infections**:

- **Dogs:** Hemoprotozoan infections (e.g., Babesia, Ehrlichia), and **von Willebrand factor** levels (if indicated)
- **Cats:** FeLV, FIV, **Haemoplasma**, and optionally **Bartonella spp.**
- **Livestock:** Should be free from **anaplasmosis**, **babesiosis**, and **theileriosis**

G. Informed Consent

- **Written informed consent** must be obtained from the owner or custodian prior to every donation.
- Donors must be **voluntary and non-remunerated** to uphold ethical standards.

H. Blood Collection and Compatibility Testing

- Blood must be collected **aseptically via jugular venipuncture** using sterile equipment.
- **Blood typing and crossmatching** are mandatory before transfusion to avoid incompatibility reactions.

I. Post-Donation Care and Observation

- All donor animals must be **monitored for at least 1–2 hours post-donation** for signs of hypovolemia or distress.
- Provide **oral fluids, light feeding, and rest**.
- Record vital parameters post-donation and advise follow-up care if needed.

7. Blood Collection and Processing

Blood collection must follow **species-specific protocols** to ensure **donor welfare**, maintain **blood quality**, and minimize transfusion risks. The **jugular vein** is the preferred site across species due to its size and accessibility. **Aseptic preparation** of the collection site is mandatory in all cases.

A. General Principles

- **Closed collection systems** are recommended for **dogs and livestock** to prevent contamination and ensure sterility.
- **Open systems** may be used for **cats**, but only when a closed system is not feasible.
- **Anticoagulants** must be used appropriately based on species and collection volume (see Table 1).
- Blood collection must be performed by **trained personnel under veterinary supervision**.

B. Blood Volume Guidelines and Anticoagulants

Species	Safe Blood Volume	Recommended Anticoagulant	Anticoagulant-to-Blood Ratio
Dogs	15–20 ml/kg body weight	CPDA-1	1 ml per 9 ml of blood
Cats	10–12 ml/kg body weight	CPDA-1 or ACD	1 ml per 9 ml (CPDA-1); 50 ml per 450 ml (ACD)
Cattle/Buffalo	Up to 1.6% of body weight (max ~6–8 L)	Sodium citrate or ACD	100 ml/L (sodium citrate); 50 ml per 450 ml (ACD)
Horses	6–8 liters	CPDA-1 or ACD	Species-specific; veterinary discretion

Note: Maximum blood volumes are based on healthy adult donors and should be adjusted based on clinical assessment.

C. Collection Protocol

- Use **sealed bags** compatible with anticoagulants for collection and storage.

- In **livestock**, use specially designed **2-liter blood collection kits** pre-loaded with anticoagulants.
- **Gentle continuous shaking** of the blood bag is necessary during collection to prevent clotting.
- Blood volume collected should be based on:
 - Donor's **body weight**
 - Expected **packed cell volume (PCV)** required for the recipient

D. Donor Monitoring and Post-Donation Care

- Monitor **vital signs** (heart rate, respiratory rate, mucous membrane color, capillary refill time) **throughout the procedure**.
- Watch for signs of **hypovolemia, weakness, or discomfort**.
- Provide **oral or IV fluid support** as appropriate post-donation.
- Observe donors for **at least 1 hour post-procedure** and record any adverse reactions.

7.1 Component Separation and Storage

Post-collection, blood may be separated into red blood cells, plasma, and platelets via centrifugation. It is important to mention expiry dates clearly and blood labelling should meet ISO9001 tracerbility norms as much as possible. Storage requirements vary for each component:

Whole blood with anticoagulant ACD, CPD or CP2D should be stored at 1–6°C and used within 21 days (ACD) 28 days (CPD, CP2D) from date of collection and **whole blood with anticoagulant CPDA-1** should be stored at 1–6°C and used within 35 days from date of collection .

Plasma should be frozen at –18°C and is viable for up to one year.

Platelets must be stored at 20–24°C with continuous agitation and used within 7 days. Using sterile equipment and proper anticoagulants is essential to preserve each component's functionality. Blood filters are routinely employed to remove any clots that may have formed during collection or storage.

7.2 Transfusion Protocols

Transfusions must be conducted under **veterinary supervision** with species-appropriate protocols to ensure safety, efficacy, and traceability. The procedure should follow a **standard checklist**, include **pre-transfusion compatibility testing**, and ensure that **each transfusion is recorded in the animal's medical records**.

A. Pre-Transfusion Requirements

- Conduct **blood typing** and **major crossmatching** to ensure compatibility.
- Maintain a **transfusion checklist**, including:
 - Patient ID, species, weight, and clinical indication
 - Donor ID and screening record
 - Blood product type and volume
 - Date and time of transfusion
- Ensure **informed owner consent** has been obtained and documented.
- Verify **storage and expiry details** of the blood product prior to use.

B. Administration Route

- Preferred route is **intravenous** via the **jugular or cephalic vein**.
- For **neonates or critical patients**, **intraosseous access** may be used when venous access is not possible.

C. Rate and Duration of Transfusion

- **Start slow:** 0.5–1 ml/kg/hr for the first 15–30 minutes to monitor for adverse reactions.
- **Dogs and Cats:** Total transfusion must be completed within **4 hours**.
- **Farm Animals:**
 - Begin gradually for the first 30 minutes.
 - Total transfusion time should not exceed **4 hours**.

D. Equipment and Asepsis

- **Blood should be warmed** to body temperature (37°C) before transfusion.
- Use **non-heparinised saline** to prime the IV line (never mix with calcium-containing fluids).
- Employ **blood administration sets with inline filters (170–260 µm)** to remove clots and debris.
- Use **extension lines** connected from the fluid pump to the catheter to maintain sterility.
- Ensure **sterile technique** during all handling of blood products.

E. Monitoring and Documentation

- Monitor vital signs every 15–30 minutes:
 - Temperature, heart rate, respiratory rate, mucous membrane color
 - Monitor for signs of transfusion reactions (e.g., fever, urticaria, tachycardia, hemolysis)
- Document the following in the **animal's medical record**:
 - Transfusion start/end time
 - Volume and type of product transfused

- Any observed reactions and action taken

7.3 Monitoring and Reaction Management

Close monitoring of the recipient animal during and after transfusion is essential for early identification of adverse reactions and assessment of transfusion effectiveness. All observations and actions must be **clearly documented in a dedicated transfusion logbook**, which serves as a critical tool for **traceability, audit, and future reference**.

A. Monitoring Schedule

- **During transfusion:**
 - **First hour:** Monitor vital signs every **15 minutes**
 - **Subsequent period:** Monitor every **30–60 minutes**
- **Post-transfusion:**
 - Evaluate **Packed Cell Volume (PCV)** 2–4 hours after transfusion to assess efficacy
 - Continue clinical observation for **at least 2 hours** post-transfusion

B. Common Signs of Acute Transfusion Reaction

- **In farm animals:** Increased respiratory rate, tachycardia, sweating, muscular tremors, violent movements
- **In dogs and cats:** Fever, vomiting, facial swelling, restlessness, panting, urticaria, or collapse

C. Emergency Response Protocol

- **Stop transfusion immediately** if any acute reaction is suspected.
- Administer appropriate **supportive care**:
 - **Antihistamines** (e.g., diphenhydramine)
 - **Corticosteroids** (as per veterinary discretion)
 - **IV fluids** if required—but **avoid hypotonic or calcium-containing solutions** immediately after transfusion due to the risk of hemolysis

D. Documentation and Traceability

- Maintain a **dedicated transfusion logbook** capturing the following:
 - Donor ID and screening details
 - Recipient information and clinical indication
 - Type and volume of blood product administered
 - Start and end time of transfusion
 - Vital signs and monitoring notes
 - Any adverse reactions and treatment administered

- This record must be appended to the **animal's medical file** and retained for institutional audit and clinical reference.

7.4 Labeling

Labeling of blood and blood components must be **accurate, standardized, and traceable** to ensure safe clinical use, quality assurance, and regulatory compliance. Each unit must carry a **tamper-proof label, wherever possible printed**, with the following mandatory details:

A. Essential Product Information

- **Product name** (e.g., Whole Blood, Packed RBCs, Plasma, Platelets)
- **Donor species** (e.g., canine, feline, bovine)
- **Blood type/group** of the donor
- **Collection date** and **expiration date**
- **Volume collected**

B. Donor and Source Identification

- **Donor ID number**
- **Unique lot or control number** (for full traceability)
- **Name and address** of the facility or manufacturer

C. Process and Additive Information

- Name and volume of **anticoagulant used** (e.g., CPDA-1, ACD)
- Details of any **additive solutions**
- **Leukoreduction status** (e.g., “Leukoreduced” if performed)

D. Storage and Handling

- **Recommended storage conditions** (e.g., 1–6°C, frozen $\leq -20^{\circ}\text{C}$)
- **Storage expiry timeline** based on the product type


E. Tube Segment Identification

- Each **associated tubing segment** (pilot tube) must bear the **same unique lot or donor ID number** to allow for re-testing or crossmatching, if required.

Labels must be **legible, permanently affixed**, and printed in English (and optionally in the local language).

8. Establishment of Blood Banks

8.1 Layout (As per TANUVAS guidelines)



8.2 Infrastructure Requirements

A functional and ethically managed veterinary blood bank must be supported by appropriate **infrastructure**, **personnel**, and **regulatory oversight** to ensure **biosafety**, **traceability**, and **clinical effectiveness**. The design and operation of the facility must adhere to **standardized layouts**, **biosafety practices**, and **quality assurance protocols**.

A. Facility Layout and Biosafety

- The physical infrastructure must follow a **standardized layout** with clearly defined zones to prevent cross-contamination:
 - **Blood collection area** (sterile environment)
 - **Component separation/processing room**
 - **Storage room** (cold chain maintenance)
 - **Record-keeping/administrative area**
- Facilities should conform to **Biosafety Level 2 (BSL-2)** practices, including:
 - Controlled access
 - Proper ventilation and airflow
 - Surface disinfection protocols
 - Use of personal protective equipment (PPE)
- A **site plan** may preferably be vetted by State Veterinary Department before operationalization.

B. Essential Equipment

- **Cold storage:**
 - Refrigerators (1–6°C) for whole blood and RBCs.
 - Freezers (–20°C or lower) for plasma.
 - Temperature loggers with alarms and backups.
- **Laboratory and transfusion support:**
 - Bench centrifuges for blood component separation.
 - Blood bag sealers, weighing scales.
 - Blood typing and crossmatching kits (species-specific).
 - Leukoreduction filters (if applicable).
 - Sterile, gas-diffusible blood collection bags with anticoagulants.
 - Continuous power supply (UPS/generators).

C. Staffing and Critical Care Access

- **24/7 access to critical care facilities** must be ensured, especially in veterinary teaching hospitals or referral centers where blood banks are established.
- Personnel must include:
 - **Veterinary transfusion-trained clinicians** to supervise donor selection, compatibility testing, and transfusions.
 - **Qualified laboratory technicians** trained in blood handling, biosafety, and quality control.
 - **Support staff** trained in asepsis, donor care, and documentation.
- Regular **training and capacity building** should be mandated for all staff involved in transfusion services.

D. Institutional Licensing and Regulatory Oversight

- Until a **national regulatory framework** is formally established, **state-level registration** of veterinary blood banks should be made mandatory.
- Blood banks must be registered with the **State Animal Husbandry Department**, subject to periodic inspections and adherence to SOPs.
- An appropriate mechanism for developing accreditation framework in the line of human blood transfusion is recommended to eventually regulate and accredit facilities nationwide.

8.3 Ideal Locations for Veterinary Blood Banks

Veterinary blood banks should be strategically located in institutions that possess the necessary veterinary infrastructure, clinical capacity, and round-the-clock emergency care facilities. The following are ideal host institutions:

- **Veterinary colleges and universities**
- **Referral hospitals and polyclinics**
- **Large veterinary diagnostic centres**
- **Government-operated multispecialty animal hospitals**

Key criteria include the presence of **transfusion-trained veterinarians**, **diagnostic laboratories**, and **critical care support**, ensuring that both donor welfare and recipient safety are maintained.

9. SOPs and Documentation

Robust Standard Operating Procedures (SOPs) and meticulous documentation form the backbone of safe and ethical transfusion practices. Institutions operating veterinary blood banks must maintain the following records:

- **Donor registration and informed consent forms**
- **Donor health screening and testing records**
- **Blood collection and labeling forms**
- **Component processing logs**
- **Inventory, storage, and expiry tracking**
- **Transfusion monitoring and reaction management forms**
- **Post-transfusion outcome assessments**
- **Waste disposal and biosafety logs**

All records must be securely maintained for a **minimum of five years**, and available for audit by regulatory or licensing authorities. Regular internal reviews should be undertaken to improve SOP adherence and address deviations.

10. Legal Framework and Regulatory Considerations

India currently lacks a dedicated national law governing animal blood banking and veterinary transfusion medicine. However, several existing statutes can provide a **regulatory foundation** for institutional governance, ethical compliance, and clinical standardization:

A. Applicable Legal Provisions

1. **Prevention of Cruelty to Animals Act, 1960**
 - Ensures donor animal welfare, humane handling, voluntary non-remunerated donation, and post-donation care.
 - Blood collection and transfusion must comply with ethical norms defined under this Act.
2. **Indian Veterinary Council Act, 1984**
 - Provides regulatory authority to the Veterinary Council of India (VCI) and State Veterinary Councils to oversee clinical veterinary practices.
 - Transfusion medicine procedures will be formally integrated into **veterinary clinical guidelines** under this Act, including licensing, training, and accreditation of transfusion centers.
3. **Biomedical Waste Management Rules, 2016**
 - Blood banks must comply with the Environment (Protection) Act provisions for disposal of biomedical waste, such as used syringes, blood bags, and contaminated materials.
4. **Drugs and Cosmetics Rules, 1945 (Indicative Human Model)**
 - While framed for human blood banks, **Part X-B** of the Rules (Rules 122-F to 122-P) offers a model structure for:
 - Licensing
 - Infrastructure standards
 - Donor safety
 - Quality control and traceability
 - These may be **adapted to veterinary needs** and act as a regulatory reference until a dedicated framework is notified.

B. Recommended Additions and Reforms

- Animal blood banks should be:
 - **Registered with the State Animal Husbandry Department** (as an interim regulatory mechanism).
 - **Eventually brought under the oversight of a national mechanism** as per the prevailing practices of human blood transfusion.
- Transfusion services be **recognized as clinical procedures** requiring licensing under the Indian Veterinary Council Act.
- Establishment of animal blood banks must ensure:
 - **Institutional accountability**
 - **Ethical donor selection**

- **Standardized infrastructure**
- **Quality assurance practices**

9. Ethical Considerations and Owner Education

Ethical principles must guide every aspect of veterinary blood donation and transfusion services. Donor animals are not mere biological resources but sentient beings whose welfare must be fully respected throughout the process. The following ethical commitments must be upheld:

- **Voluntary, non-remunerated donation** must be the cornerstone of veterinary blood banking. No monetary incentives should be offered to pet owners or livestock keepers.
- **Regular health checks** are essential to ensure donor suitability and welfare, with clear intervals between donations.
- **Humane handling, rest, and post-donation care** must be guaranteed for every donor animal.
- **Mandatory owner consent** (informed and written) should be obtained prior to every donation.
- **Awareness campaigns**, such as the proposed “**Gift of Life**” initiative, should be launched to promote community participation and build a sustainable and ethical donor base.

Pet owners and farmers must be made aware that donating blood can help save another animal's life and contribute to a humane and responsive veterinary care system.

10. National Veterinary Blood Bank Network (N-VBBN)

To facilitate standardized and equitable access to veterinary blood products across India, a **National Veterinary Blood Bank Network (N-VBBN)** may be considered under the Department of Animal Husbandry & Dairying (DAHD), Government of India. The network would serve as a national platform for:

- **Digitally integrated donor registries**, categorized by species, breed, location, and blood type.
- A **real-time inventory management system**, mapping availability of whole blood and components across participating centers.
- A **helpline and online portal** to connect veterinary clinics, hospitals, and donors in emergency situations.
- **Standardization of practices**, reporting formats, and adverse reaction logs across all registered blood banks.

Such a network would not only improve logistical efficiency but also promote transparency, traceability, and quality control across the country.

11. Training and Curriculum Development

Capacity-building is critical to the successful implementation of veterinary blood banking systems. Key recommendations include:

- **Integration of transfusion medicine** topics into the **Bachelor of Veterinary Science & Animal Husbandry (BVSc & AH)** curriculum, with emphasis on donor selection, transfusion protocols, and reaction management.
- **Postgraduate (MVSc)** courses should offer specialized modules on transfusion science and clinical transfusion medicine.
- **Continuing Veterinary Education (CVE)** programs must be developed for practicing veterinarians, including hands-on workshops and online certification courses.
- **Nodal training institutes** can be identified to act as centers of excellence in transfusion medicine, supporting regional knowledge dissemination.

12. Future Directions

To align with evolving clinical needs and technological advancements, the veterinary transfusion sector must pursue the following innovations:

- Development of **mobile applications** for donor-recipient matching, scheduling, and logistics support.
- Establishment of **cryopreservation and biobanking units** to extend storage life and support rare blood type availability.
- Research into **autologous transfusions** (self-donation) and **xenotransfusion techniques**, where feasible and ethically permissible.
- Introduction of **mobile blood collection units** to expand access in underserved rural and peri-urban areas.
- Regular **impact assessments** and feedback loops to refine practices and incorporate new findings.

References

1. Standard Operating Procedures for Veterinary Blood Banking and Transfusion by TANUVAS Animal Blood Bank, Department of Veterinary Clinical Medicine
2. Blood Transfusion in Animals: A review by Aravind S and Ninan Jacob
3. California Animal Blood Banking Guidance Resource by California Department of Food and Agriculture